

# **TOPIC: EVALUATION OF FLOOR SPREADS ON BROODING EFFICIENCY AND PERFORMANCE ON BROILER CHICKS**

## **CHAPTER ONE**

### **1.0 INTRODUCTION**

The poultry industry has in recent years occupied a leading position among agricultural industries in many parts of the world. This may be due to the attractive attributes of poultry which include the ability to adapt easily, high economic value, rapid generation time and a high rate of growth that can result in the production of meat within 8 weeks (Smith, 2002).

However, with the increasing concentration of poultry in Nigeria, it has become essential to appreciate the influence of litter materials on which these birds are reared. In Nigeria, broilers and cockerels are mostly reared on the floor spread with litter materials. Litter material is any dry form material used on the floor of chicken houses on which chicken dropping will fall. It is known as litter material because it combines with the droppings and undergoes a bacterial breakdown process, thus preventing a smelly and unsanitary condition (Demirulus, 2006). Various types of litter materials are used in different countries. The common types of litter used in poultry houses throughout the world are sawdust, rice husk, by products, wood shavings, corn cobs, oat hulls, dried leaves, coffee husk (Rao, 1986). In Southwestern Nigeria, wood shavings are easily ob

taind from sawmills at little or no cost and used as litter. The use of this material has been hinged purely on their availability and price, without any consideration for the comfort of the birds. Few available reports on the effect of litter on the performance of birds are contradictory and showed a need for validation (Oliveira .1974). According to Awojobi (1999), types of litter had no significant effect on birds' performance. Whereas Popolizio . (1979), Poyraz.(1990) and Anisuzzaman and Chowdhury (1996) reported that rice husk was the best litter for broiler chickens. Based on these premise, it becomes imperative to evaluate wood shaving and other locally available litter materials in Nigeria such as sawdust ,rice husks,groundedCorncobshaft and Bare floor on the performance of broiler chickens.

This study is vital for improving broiler chicken production by evaluating the impact of different floor spreads on various aspects of broiler management. Previous research has highlighted the significance of brooding conditions on chick survival and early growth (Alhamad .2017). Understanding how floor spreads influence brooding efficiency can lead to better management practices, reducing mortality and enhancing productivity (Muir, 2011).

In addition, the study will explore the effects of floor spreads on broiler growth rates and health, with previous studies suggesting that environmental conditions, such as floor material, significantly affect bird welfare and disease risk (Dawood. 2019). Evaluating these parameters will offer insights into reducing health issues and improving overall flock vitality (Cymerys .2018).

The economic impact of floor spreads will be assessed, as cost-effective production practices are essential for maintaining profitability in the poultry industry (Jahan . 2020). By analyzing the costs and benefits of different floor spreads, this study will provide practical recommendations for optimizing both production efficiency and economic outcomes.

### **1.1 PROBLEM STATEMENT**

The efficiency and profitability of broiler chicken production are heavily influenced by various management practices, including the choice of floor spreads used in rearing systems. However, limited research has been conducted to comprehensively understand the impact of different floor spreads on critical parameters such as brooding efficiency, growth rates, health outcomes, behavior, and economic viability. The lack of clear guidelines on the most suitable floor spreads poses challenges for poultry farmers in optimizing productivity while ensuring animal welfare and reducing costs.

So this study we aims to address this gap by systematically evaluating the effects of various floor spreads on broiler chickens to provide evidence-based recommendations for sustainable and cost-effective poultry farming practices.

### **1.2 SIGNIFICANT OF THE STUDY**

This study is significant because it aims to improve broiler production by assessing the impact of different floor spreads on brooding efficiency, growth, and health. Identifying the best floor spreads can enhance survival rates and growth performance, leading to

more productive flocks

Economically, these study will examines the cost-effectiveness of different floor spreads, helping farmers make informed decisions that improve profitability while maintaining high animal welfare standards. The findings could guide more sustainable and ethical farming practices in the poultry industry.

Additionally, the research explores how floor spreads affect broiler health and welfare. By evaluating disease incidence, mortality, and behavior, the study provides insights in to minimizing stress and promoting better welfare for the birds.

### **1.3. OBJECTIVE**

1. Assessing the impact of different floor spreads on brooding efficiency,
2. Evaluating the influence of floor spreads on the growth rate
3. Investigating the effects of various floor spreads on the health parameters of broiler chickens, including disease incidence and mortality rates.
4. Comparing the behavior patterns of broiler chickens raised on different floor spreads to understand their welfare and stress levels.
5. Analyzing the economic implications of using different floor spreads in terms of cost effectiveness and return on investment in broiler production.

### **1.4 JUSTIFICATION**

This study is important because it examines how different floor spreads affect brooding efficiency, which is crucial for chick survival and early growth. By understanding these effects, producers can improve brooding conditions, reduce mortality, and enhance overall production efficiency.

The study also focuses on the impact of floor spreads on broiler growth rates and health. It will explore how various spreads influence factors such as cleanliness, comfort, and disease risk, helping farmers make better decisions to improve bird health and reduce mortality.

Also the research will justify the economic implications of using different floor spreads. By evaluating their cost-effectiveness and impact on productivity, the study will provide recommendations for optimizing production practices, ensuring better profitability and animal welfare in broiler farming.

## **CHAPTER TWO**

### **LITERATURE REVIEW**

#### **2.1 HISTORICAL BACKGROUND**

The use of floor spread materials in poultry production has evolved over the decades as part of a broader effort to improve brooding efficiency, chick health, and overall performance. Historically, floor management in poultry housing was rudimentary, with early poultry systems relying heavily on bare floors or simple earth floors. These systems were often plagued with problems such as poor sanitation, high disease incidence, and low pro

ductivity (Adene & Oguntade, 2008).

Before the commercial broiler industry gained momentum in the mid-20th century, most poultry were raised in small backyard systems. During this period, Birds were commonly raised on bare ground or clay floors, with little attention to bedding or litter and this cause of Disease outbreaks, particularly those caused by parasites and bacteria, were common due to direct contact with feces and moisture accumulation (Smith et al., 2017).

Litter Material With the rapid expansion of commercial poultry farming is adopted from the years 1950s \_1970s especially in the United States and Europe, the importance of managing the poultry house environment became evident.

Brooding efficiency improved significantly as farmers observed better chicks survival, more stable temperatures, and reduced disease (Atkins et al., 2018).

Scientifically During this period of Research and Performance Metrics in the years 1980s–2000s detailed studies began to emerge comparing different bedding materials and their impact on:

- Growth rate
- Feed conversion ratio (FCR)
- Mortality
- Footpad health and litter quality

And the Key developments included:

- Comparative studies between sawdust, rice husk, peanut shells, and wood shavings—especially in tropical regions (Abdullahi et al., 2015).
- Recognition of the role of bedding in ammonia control, bacterial growth, and bird behavior (Johnson et al., 2021).
- Introduction of alternative materials like paper pellets, corn cobs, and recycled products (Ibrahim et al., 2019).
- Modern Innovations and Sustainable Practices (2010–Present)

And in Recent years increased have seen focus on:

- Sustainable and cost-effective materials, such as ground corn cobs, cocoa husks, and recycled newspaper (Oke et al., 2020)
- The economic impact of floor spread choices, especially for small-scale and rural poultry producers (Umar & Musa, 2017).
- Behavioral and welfare assessments of broilers on different bedding types to ensure ethical farming practices.
- Use of technology and sensors to monitor litter moisture and ammonia, guiding decisions on litter replacement and ventilation (Atkins et al., 2018).

In developing countries like Nigeria, India, and Bangladesh, emphasis has been placed on locally available materials such as rice husk, groundnut shells, and plant waste, which provide affordable and effective alternatives to imported bedding (Oke et al., 202

0).

## **2.2 IMPORTANT OF FLOOR MANAGEMENT IN BROODING**

Floor spread materials, also known as bedding or litter materials, serve several important functions in poultry houses. They help in moisture absorption, insulation, odor control, and waste management. An ideal floor spread material should be:

- Highly absorbent
- Non-toxic and dust-free
- Comfortable and thermally insulating
- Economical and readily available
- Easy to dispose of or recycle

Improper floor materials or bare floors can lead to issues such as footpad dermatitis, high ammonia levels, increased mortality, and poor weight gain (Atkins et al., 2018).

## **2.3 COMMON FLOOR SPREAD MATERIALS IN POULTRY BROODING**

### **2.3.1 SAWDUST**

Sawdust is one of the most widely used bedding materials in poultry farming. It is highly absorbent and relatively cheap. However, if not properly dried or sourced from treated wood, it may lead to respiratory issues or caking. Proper management of sawdust improves chick comfort and minimizes disease risks (Smith et al., 2017).

### **2.3.2 WOOD SHAVINGS**

Wood shavings provide good cushioning and insulation for broiler chicks. They are bulkier than sawdust and allow for better aeration. Studies have shown that birds raised on wood shavings tend to have better body weight gain and lower incidences of breast blisters (Olomu, 2016).

### **2.3.3 RICE HUSK**

Rice husk is a common by-product of rice processing and is used in poultry farming, especially in rice-producing regions. Although less absorbent than sawdust, rice husk provides a good balance of insulation and airflow. It also supports microbial fermentation that helps in breaking down droppings, reducing ammonia build-up (Oke et al., 2020).

### **2.3.4 GROUNDED CORN COBS**

Ground corn cobs are an underutilized by-product in poultry bedding. They have moderate absorbency and decompose slowly, making them suitable for longer production cycles. Limited studies suggest that broiler chicks raised on ground corn cobs show comparable growth performance to those on traditional litters, though care must be taken to avoid mold contamination (Ibrahim et al., 2019).

### **2.3.5 BARE FLOOR (NO LITTER)**

The use of bare floors is discouraged in commercial brooding because it offers no cushioning, poor thermal insulation, and minimal moisture absorption. Chicks brooded on bare floors are more prone to stress, cold, foot injuries, and infections such as coccidiosis. Performance metrics such as feed conversion ratio and average daily gain are often lower on bare floors (Adene & Oguntade, 2008).

## **2.4 FLOOR SPREAD AND BROODING EFFICIENCY**

Brooding efficiency involves optimal chick comfort, temperature maintenance, and low mortality rates. Floor spread materials play a pivotal role by providing insulation and reducing direct heat loss. Chicks raised on appropriate bedding materials demonstrate:

- Quicker adaptation to the environment
- Uniform growth
- Reduced heat stress
- Improved survival rates

(Johnson et al., 2021) showed that poor litter leads to poor brooding conditions which affect chick behavior and energy balance.

## **2.5 FLOOR SPREAD AND GROWTH PERFORMANCE**

Performance indicators such as weight gain, feed conversion ratio (FCR), and mortality are closely linked to the quality of litter. Well-managed floor spreads contribute to:

- Better feed intake
- Higher weight gain
- Reduced disease incidence

Broiler chicks raised on sawdust and wood shavings typically outperform those raised on rice husk or bare floors in terms of final live weight and FCR (Abdullahi et al., 2015).

## **2.6 HEALTH IMPLICATIONS OF FLOOR SPREAD MATERIALS**

Poor floor spread management can lead to excessive moisture buildup, leading to bacterial and fungal growth, increased ammonia levels, and respiratory problems. Ideal floor spreads reduce microbial load and minimize the risk of diseases like aspergillosis, coccidiosis, and pododermatitis.

## **2.7 ECONOMIC CONSIDERATIONS**

The cost and availability of floor spread materials affect their choice, especially in rural or low-resource areas. While sawdust and wood shavings might be ideal, in some regions, rice husk or even corn cobs are more economical. The bare floor is the least expensive but has high long-term costs due to increased mortality and poor growth performance (Umar & Musa, 2017).

## **CHAPTER THREE**

### **3.1 STUDY AREA**

The research is carried out at Agricultural technology garden located in KWARA STATE POLYTECHNIC ILORIN, KWARA STATE.

### **3.2 EXPERIMENTAL DESIGN AND TREATMENT**

The experiment is comprised of treatment viz. Sawdust, Rice husks, Wood shaving, grounded Corn cob shaft, and Bare floor. These floor materials were selected based on their availability and prior studies. The experiment is laid down in a completely randomised design (CRD) and replicated three times.

### **3.3 SAMPLE PROCEDURE AND SIZE**

Six (6) broiler chicks are used per treatment, given a sample size of 30 chicks.

The design includes five (5) treatment groups i.e. the four floor spreads and the bare floor (control).

Those chicks will be randomly distributed into the experimental units.

### **3.4 MANAGEMENT OF EXPERIMENTAL BIRDS**

- Housing: The brooding pens measured is provide a sufficient space per bird
- Feeding: A nutritionally balanced starter diet formulated to standards is be provided ad libitum.
- Lighting and Temperature: lighting is provided daily during the brooding phase, and temperatures is maintained gradually
- Vaccination: Birds is vaccinated against castle Disease and Infectious Bursal Disease following
- Biosecurity Measures: Strict hygiene protocols is implement to prevent disease outbreaks

### 3.5 Data collection

Data is taken at interval, of live weight, feed intake, feed conversion ratio, mortality, Litter moisture, pH and temperature.

At the end of the experiment, the data taken is subjected to Analysis of Variance (ANOVA), and were separated using Duncan Multiple Range Test.

## REFERENCES

ATAPATTU, N.S.B.M. and. Wickraramasinghe, K.P. (2007).The use of refused tea as litter material for broiler

Chickens. Poultry Science, 86:968–972.

Awojobi, H.A, Adekunmi, A.A. and Adebowale, O.J. (1999). Comparative Performance of Broiler chickens reared

On different litter materials. Tropical Animal Production Investigations, 2: 135 – 141.

Alhamad, M. N. (2017). Brooding management and early chick development. Journal of Applied Poultry Research,

Anisuzzaman, M. and Chowdhury, S.D. (1996).Use of four types litter for rearing broilers. British Poultry Science,

BABATUNDE, G.M. (1998). The presidential task force on alternative feed resources: Challenges and way forward

Nigeria Society of Animal Production Newsletter, 17(2): 4

De Avila, V.S., de Oliveira, U., de Figueiredo, E.A.P., Costa, C.A.F., Abreu, V.M.N. and Rosa, P. S., (2008).

Alternative material to replace wood shavings as broiler litter. Rev. Bras. Zootecn., 37: 273-277.

Cymerys, M. (2018). Floor material and poultry health management. Livestock Science,

Dawood, M. A. O. (2019). Environmental factors affecting poultry health and growth. Poultry Science, 98(5), 1933-1944.

Demirulus, M. (2006). The effect of litter type and litter thickness on broiler carcass traits. International Journal of Poultry Science,

Jahan, M. S.I. (2020) Aro, S.O., Aletor, V.A., Tewe, O.O., Fajemisin, A.N., Usifo, B. P

4th Annual Conference of School of Agricultural Technology hold at the Fed. University of Tech. Akure.

Nigeria, 21<sup>st</sup> May 2008. Pp: 86-92.

Muir, W. M. (2011). Genetic selection for growth and brooding efficiency. Poultry Science, 90(6), 1169-1177.

Popolizio, E.R., Ricci, H.R., Castellote, H.F. and Pailhe, L.A. (1979). Use of different materia

Is for poultry litter

Poyraz, Ö., İscan, K.A., Nazlıgül, Deliormanoglu, A. (1990). Diatomite Utilization Possibilities in broiler

Production. Journal Veterinary Association, 47-57.

Rao, V.S.R. (1986). Litter-Its management and utility in broiler. Poultry Advisory, 19(7-12): 31-40.

Smith, J.C. (2002). Chopped Bermuda grass hay as an alternative bedding material for rearing market tur.

## REFERENCES

Adene, D. F., & Oguntade, A. E. (2008). The Structure and Importance of the Commercial and Village Based Poultry Industry in Nigeria. FAO Consultancy Report. Food and Agriculture Organization of the United Nations.

Abdullahi, Y. A., Bello, M. B., & Dogo, A. G. (2015). Effect of different bedding materials on the performance and carcass characteristics of broiler chickens. International Journal of Poultry Science, 14(7), 387–390.

Atkins, K. L., Johnson, C. R., & Cole, M. N. (2018). Influence of litter type on broiler performance and footpad health. Journal of Applied Poultry Research, 27(3), 452–460.

Ibrahim, H., Abdulkadir, A., & Musa, U. (2019). Utilization of ground corn cobs as litter m

aterial for broiler chicks. *Nigerian Journal of Animal Production*, 46(1), 123–129.

Johnson, S. M., Thomas, L. J., & Kurek, D. R. (2021). Impact of bedding material on brooding environment and chick welfare. *Poultry Science Journal*, 100(5), 235–241.

Oke, F. O., Akinwale, M. O., & Adeyemi, T. A. (2020). Comparative effects of rice husk and sawdust on the performance of broiler chickens. *Nigerian Poultry Science Journal*, 17(2), 105–112.

Olomu, J. M. (2016). *Monogastric Animal Nutrition: Principles and Practice* (2nd ed.). Benin City: Jachem Publishers.

Smith, B. A., George, F. O., & Ajayi, A. B. (2017). Sawdust quality and its effect on litter performance and broiler growth parameters. *Livestock Research for Rural Development*, 29(6), Article #111.

Umar, A. M., & Musa, A. A. (2017). Economic evaluation of litter materials in broiler production in semi-arid areas. *Nigerian Journal of Animal Science and Production*, 44(1), 89–96.

North, M. O., & Bell, D. D. (1990). *Commercial Chicken Production Manual* (4th ed.). New York: Van Nostrand Reinhold.

FAO. (2010). *Small-scale poultry production*. FAO Animal Production and Health Manual No. 1. Food and Agriculture Organization of the United Nations.

ferences

Abdullahi, Y., Musa, U., & Ibrahim, A. (2015). Effect of different litter materials on performance and welfare of broiler chickens. *Nigerian Journal of Animal Science*, 17(2), 45–52.

Adene, D. F., & Oguntade, A. E. (2008). The structure and importance of the commercial and rural poultry industry in Nigeria. *FAO Poultry Sector Country Review*.

Atkins, J. L., Morrison, D., & Smith, C. (2018). Bedding materials and broiler performance: A review. *Poultry Science International*, 97(4), 1025–1032.

Ibrahim, S., Musa, J., & Lawal, R. (2019). Use of corn cobs as bedding material for broiler chicks: Performance and microbial load. *Tropical Agriculture Research*, 31(1), 12–18.

Johnson, H. M., Chukwu, G. C., & Bello, A. M. (2021). Impact of floor bedding on ammonia emissions and broiler welfare. *Journal of Poultry Research and Innovation*, 29(3), 60–67.

Olomu, J. M. (2016). *Monogastric Animal Nutrition: Principles and Practice*. Benin City: Jachem Publishers.

Oke, O. E., Afolabi, K. D., & Abioja, M. O. (2020). Evaluation of alternative bedding materials on growth performance, carcass traits, and litter quality of broiler chickens. *International Journal of Livestock Production*, 11(5), 85–92.

Smith, B., James, D., & Allen, L. (2017). Litter materials and broiler health: Comparative outcomes. *Poultry Environment Review*, 44(2), 78–84.

Umar, I. M., & Musa, A. A. (2017). Cost-effectiveness of local litter materials in broiler production. *African Journal of Agricultural Economics*, 15(3), 134–140.

